

1. Niggemann B, Lange L, Finger A, Ziegert M, Müller V, Beyer K, Accurate oral food challenge requires a cumulative dose on a subsequent day., *J. Allergy Clin. Immunol.* 2012;130(1): 261–263.
2. Trendelenburg, V., Ahrens, B., Wehrmann, A.-K., Kalb, B., Niggemann, B., Beyer, K. Peanut allergen in house dust of eating area and bed - a risk factor for peanut sensitization? *Allergy* 2013; 68(11):1460–1462.
3. Blumchen K., Beder A., Beschorner J., Ahrens F., Gruebl A., Hamelmann E., Hansen G., Heinzmann A., Nemat K., Niggemann B., Wahn U., Beyer K. Modified oral food challenge used with sensitization biomarkers provides more real-life clinical thresholds for peanut allergy. *The Journal of Allergy and Clinical Immunology* 2014; 134(2):390-8.
4. Beyer K., Grabenhenrich, L., Beder, A., Kalb, B., Ziegert, M., Finger, A., et al. (2014). Predictive values of component-specific IgE for the outcome of peanut and hazelnut food challenges in children. *Allergy* 2015; 70: 90–98.
5. Trendelenburg, V., Enzian, N., Bellach, J., Schnadt, S., Niggemann, B., & Beyer, K. Detection of relevant amounts of cow's milk protein in non-pre-packed bakery products sold as cow's milk-free. *Allergy* 70(5):591-7.
6. Grabenhenrich L, Lange L, Härtl M, Kalb B, Ziegert M, Finger A, Harandi N, Schlags R, Gappa M, Puzzo L, Stephan V, Heigele T, Büsing S, Ott H, Niggemann B, Beyer K. The component-specific to total IgE ratios do not improve peanut and hazelnut allergy diagnoses. *J Allergy Clin Immunol.* 2016 Jun;137(6):1751-1760.
7. Schoemaker AA, Sprickelman AB, Grimshaw KE, Roberts G, Grabenhenrich L, Rosenfeld L, Siegert S, Dubakiene R, Rudzeviciene O, Reche M, Fiandor A, Papadopoulos NG, Malamitsi-Puchner A, Fiocchi A, Dahdah L, Sigurdardottir ST, Clausen M, Stańczyk-Przyłuska A, Zeman K, Mills E, McBride D, Keil T, Beyer K. Incidence and natural history of challenge-proven cow's milk allergy in European children -EuroPrevall birth cohort. *Allergy.* 2015 Aug;70(8):963-72.
8. Xepapadaki P, Fiocchi A, Grabenhenrich L, Roberts G, Grimshaw KE, Fiandor A, Larco JI, Sigurdardottir S, Clausen M, Papadopoulos NG, Dahdah L, Mackie A, Sprickelman AB, Schoemaker AA, Dubakiene R, Butiene I, Kowalski ML, Zeman K, Gavrili S, Keil T, Beyer K. Incidence and natural history of hen's egg allergy in the first 2 years of life-the EuroPrevall birth cohort study. *Allergy.* 2016 Mar;71(3):350-7.
9. Lange L, Lasota L, Finger A, Vlajnic D, Büsing S, Meister J, Broekaert I, Pfannenstiel C, Friedrichs F, Price M, Trendelenburg V, Niggemann B, Beyer K. Ana o 3-specific IgE is a good predictor for clinically relevant cashew allergy in children. *Allergy.* 2017 Apr;72(4):598-603
10. Bellach J, Schwarz V, Ahrens B, Trendelenburg V, Aksünger Ö, Kalb B, Niggemann B, Keil T, Beyer K. Randomized placebo-controlled trial of hen's egg consumption for primary prevention in infants. *J Allergy Clin Immunol.* 2017 Epub ahead of print